

## Impact of Automobile Emissions on the Productivity of *Crocus sativus* L.

Rafiq, S. K.<sup>1\*</sup>, Ganai, B. A.<sup>2</sup> and Bhat, G. A.<sup>1</sup>

<sup>1</sup>Department of Environmental science, University of Kashmir, Hazratbal, Srinagar  
Jammu & Kashmir 190006, India

<sup>2</sup> Department of Biochemistry, University of Kashmir, Hazratbal, Srinagar Jammu &  
Kashmir 190006, India

Received 27 Sep 2007;

Revised 10 Feb 2008;

Accepted 25 Feb 2008

**ABSTRACT:** Effect of automobile pollution on the productivity of *Crocus sativus* was studied for one year time period (October 2005-April 2006). The parameters which were analyzed viz total chlorophyll, carotenoids, leaf area, plant height, corm size, corm weight, corm yield, biomass, flower production, and length of stigma and style, showed a remarkable decrease as compared to control. Total chlorophyll, carotenoids and leaf area decreased by approximately 40%, plant height by 36%, fresh biomass by 33%, dried biomass by 25%, corm yield by 29%, corm number by 32%, corm size by 33%, flower production by 33%, length of stigma+style by 23%, fresh weight of flower by 15% and dried weight of flower by 14% respectively.

**Key words:** Automobile emissions, Productivity, *Crocus sativus*, Suspended particulate matter, SO<sub>2</sub>, NO<sub>2</sub>

### INTRODUCTION

Saffron (*Crocus sativus*.) is highly prized as one of the best natural food flavorings and coloring substance having general panacea properties. Chief pigment of saffron is its yellowish red glycoside crocin. *Crocus sativus* is a perennial low growing herb with globular corms ranging from 0.5 to 5.0 cm in diameter. The sizes of the corm have a profound effect on the productivity of *Crocus sativus* (Ingram 1969, Madan *et al.* 1996). Besides corm size saffron production is also influenced by photoperiod, humidity, atmospheric composition, etc. (Madan *et al.*, 1996). The atmospheric composition is altered by human activity like air pollution. As per estimated automobile emissions is the primary source of air pollution (Sivasamy and Srinivasan, 1997). The polluted air may have profound effect on the growth and development on vegetation, because gases enter the leaves during the normal course of gaseous exchange and cause blocking of

stomata, chlorophyll degradation, less starch synthesis, damage to leaves and other tissues, growth reduction and ultimately decline in crop yield (Grime 1970, Well burn *et al.*, 1981, Bates and Farmer, 1992, Agarwal *et al.*, 2003, and Yi *et al.*, 2005).

### MATERIALS & METHODS

The entitled work was carried out from October 2005 to April 2006. Pampore karewa (Jammu & Kashmir, India), where it is cultivated as a cash crop, was selected as a site to study the impact of auto emission on saffron productivity. The crop production in this region had faced a decline from past few years. One of the factors which have been associated with it is the chronic exposures of *Crocus sativus* to automobile emissions. Following are the methods which were adopted to monitor air quality of the region and to analyze different plant parameters High volume air sampler was used to monitor NO<sub>2</sub>, SO<sub>2</sub> and

\*Corresponding author E-mail: skrafiq@earthling.net

suspended particulate matter .NO<sub>2</sub> estimation was done by Merryman *et al.* (1973) method, SO<sub>2</sub> was estimated by West and Gaek (1956) method, and suspended particulate matter was estimated by filtration method. Chlorophyll and carotenoids were extracted in 80% acetone and readings were measured at 645,663,510 and 480 nm and calculations were made according to Arnon (1949) using absorption coefficient. Leaf area was measured by using a long graph paper. The plant height was measured as the maximum length of aerial shoots arising from the apical notes of a corm from the soil surface to its apex. Corms were dug from each quadrat of 1m x 1m. After removing the soil from the corms they were weighed and recordings were made. The number of corms were also counted and noted down. The dug out corms from a given quadrat of 1m x 1m were taken out, cleaned and later measured for its diameter with the help of a fiberglass tape. Number of flowers was estimated by harvesting the flowers present in an 1m x 1m quadrat. The stigmas and styles were then shaved off and measured for its length using a centimeter scale. The flowers after weighed for fresh masses were then dried in sunlight until uniform dried weight was obtained.


**RESULTS & DISCUSSION**

The effect of automobile emission on the productivity of *Crocus sativus* is summarized in Tables 1, 2 and 3 and Figs. 1-8. The results of decline in average annual productivity due to automobile emissions at various sites along roadsides are in agreement with each other. The factors most likely to influence the productivity were the synergetic effects of NO<sub>2</sub>, SO<sub>2</sub> and SPM (Table 1) and the winter conditions that prevailed during winter times, this is in agreement with the finding of Asheden and Mansfield (1978), Murray *et al.* (1992) and Bell and Treshow (2002). The effect of auto emission (NO<sub>2</sub>, SO<sub>2</sub> & SPM) had a detrimental effect on the amount of total chlorophyll and carotenoid as shown in Table 2. Total chlorophyll & carotenoids showed a declining trend along road rides as compared to control. The results are in conformity with the findings of Neubert *et al.* (1993), Tingey *et al.*(1971), Legge *et al.*(1998), Pratibha & Madhu (2000), Hao *et al.*(2002). Agarwal *et al.*(2003) and Yi, *et al.*,2005.


The data (Table 2) reveals that automobile emission also had a detrimental influence on the corm size, corm number, and corm yield along roadside. The declined trend along road sides as compared to control is attributed to the fact that the *Crocus sativus* being a perennial herb is susceptible to pollution and shows reduced photosynthetic activity and less storage of starch in corm. This is in agreement to the results of Well burn *et al.*(1981), Frevert (1984), Modi *et al.*(1992), Legge (1996), Vorobeichik (2002) and Hao *et al.* (2002).

**Table 1. Ambient air monitoring along roadside From October 2005-April 2006**

SNO	Parameters	Permissible limits Mean values	Recorded concentration
1.	SO <sub>2</sub> (µg/m <sup>3</sup> )	30	58
2.	NO <sub>2</sub> (µg/m <sup>3</sup> )	30	84
3.	SPM (µg/m <sup>3</sup> )	100	346



**Fig. 1. Comparative ambient air quality**



**Fig. 2. Concentrations of Chlorophyll and Carotenoids**




Fig. 3. Leaf and Plant characteristics




Fig. 4. Fresh and Dry biomass




Fig. 5. Corn yield and number





Fig. 6. Corn size

Table 2. Vegetative plant analysis of *Crocus sativus* from October 2005-April 2006


S. No	Parameter	Control	Exposed Sites along road sides		
			Site A	Site B	Site C
		Mean	Mean	Mean	Mean
1	T. Chlorophyll (mg/g)	1.20	0.71	0.74	0.68
2.	Carotenoids (mg/g)	0.60	0.37	0.38	0.34
3.	Leaf Area (cm <sup>2</sup> )	9.06	5.74	5.10	5.94
4.	Plant Height (cm)	22.54	14.40	14.50	15.05
5.	Fresh Biomass above ground (g/m <sup>2</sup> )	46.85	31.85	30.80	32.57
6.	Dried Biomass above ground (g/m <sup>2</sup> )	9.68	7.6	7.4	7.05
7.	Corm yield (g/m <sup>2</sup> )	355.71	252.85	248.57	261.57
8.	Corm number (NO/m <sup>2</sup> )	62.28	43.4	42.28	42.28
9.	Corn size (cm)	2.72	1.82	1.86	1.81

**Table 3. Flower production /m<sup>2</sup>, length stigma + style, fresh weight of Flowers/m<sup>2</sup> and dried weight of flowers/ m<sup>2</sup> (October and November 2005)**

S. No	Parameter	Control	Exposed Sites along road sides		
			Site A	Site B	Site C
		Mean	Mean	Mean	Mean
1	Flower Production (NO /m <sup>2</sup> )	34	23.0	22.0	24.0
2	Length of Stigma +Style (cm)	3.96	3.0	3.2	3.0
3	Fresh weight of flowers (g/m <sup>2</sup> )	3.270	2.800	2.850	2.800
4	Dried weight of flowers (g/m <sup>2</sup> )	0.483	0.417	0.421	0.414



**Fig. 7. Flower production at control and exposed sites**



**Fig. 8. Length of stigma plus style**

## CONCLUSION

Plant height, leaf area and biomass is influenced by corm size (Srinivastava, 1963,1964). During the period of study, plant height, leaf area and biomass showed a remarkable declined. These results are in total agreement with the reports of Grime (1970), Pandey (1974), Gilani (1977), Jacobson (1987), Bates & Farmer (1992), Kamalakar (1992) and Agarwal *et al.*(2003).The number of flowers / quadrat and length of stigma + style were highly influenced by auto emissions along roadside. The number of flowers and length of stigma + style is directly related to the corm size and yield. Hence the results obtained are in close proximity with that of Madan *et al.*(1966), Pandey *et al.*(1974), and Munshi and Zargar (1991).

## ACKNOWLEDGEMENTS

Our grateful thanks are due to Dr. A.R Yousuf, Head Department of Environmental Science for inspiring guidance and valuable suggestions.

## REFERENCES

- Agarwal, M., Singh, B. Rajput, M. Marshal, F. and Bell, J. N. (2003) Effect of Air pollution on peri urban agriculture a case study. Environ. Pollut., **126**(3), 323-329.
- Arnon, D.I. (1949) Acetone method for the extraction of chlorophyll from leaf tissues: Plant Physio., **24**, 1-15.
- Asheden, T.W. and Mans field T. A. (1978) Extreme Pollution sensitivity of grasses when SO<sub>2</sub> and NO<sub>2</sub> are present in the atmosphere together. Nature, **273**, 142-143.
- Bates, J. W. and farmer, A. M.(1992) Brophytes and Lichene in a changing environment. Clare don press. Oxford.
- Bell J. N. B and Treshow, M. (2002). Air Pollution & Plant Life 2<sup>nd</sup> editor. Publication. John Willy and sons, New York, U.S.A.
- Gillani, A. G. (1977) Methods of increasing saffron productivity. Temat Sb TR Azerb N.H Sadovodatu Vinogradarotiva Subtrop. Kultur (1976) 9 104 -108 (C.F) DIALOGY Information service Inc).
- Grime, J. P. (1970). People and plants in Derbyshire. Naturalist trust matlock.
- Hao, J., Li, J., Duan, L. He, K. and Dai, W. (2002) Evaluation of forest damage cost from emission:a case study in Hunan Province, China, Huan Jing Ke Xue, **23** (6), 1-5.
- Ingram, J. S. (1969). Saffron. Tropical science. 177-184.
- Jacobson J. S., Os melorki, J., Yamada, K. and Heller, I.(1987) The influence of stimulated acid rains on vegetation and reproductive tissue of cucumber (*Cucumis sativus L.*) New physio., **105**, 139-147.
- Kamalakar, J. A.(1992) Response of plants to auto exhaust pollution. Acta Bot. Indica, **20**, 84-88.
- Legge , A. H., Noral, M. and Krupa, S. V. (1996) Modeling the numerical relationship between chronic ambient sulphur dioxide exposure and tree growth. Can. J. Forest Res., **26**, 689-695.
- Legge A. H., Jagar, H. J. and Krupa, S. V. (1998). SO<sub>2</sub> in recognition of air pollution injury to Vegetation. In R.B Flagler, Ed a Pictorial atlas 2<sup>nd</sup> edition PP 33-42. Air and waste management association Pittsburg.
- Madan C.I., Kapur, B.M. and Gupta, U.S. (1966). Saffron Econom. Botan., **20** (4), 377-385.
- Merryman, E.L., Spicer, C.W. and Levy, A.(1973) Evaluation of arsinite modified Jacobs Hochheiser procedure. Environ. Sci. Tech., **7**, 1056-1059.
- Modi, S., Siddiqui, N., Rai, S. and Rai, P.K. (1992) *Foliar injuries in Diospyros Melanoxlon roxb.* Corm exposed to auto exhaust Pollution. Indian J. Appl. Bio., **7** (1), 29-32.
- Munishi, A. M. and Zargar G. H. (1991) Variation in the natural population of saffron(*Crocus sativus*) crop in Kashmir and performance of some selected sub population. Phytobredon, **7** (1&2), 62 -70.
- Murray, F., Nilson, S. and Monk, R. (1992) NO<sub>2</sub> & SO<sub>2</sub> mixture stimulates barley grain production but depress clover growth. Environ. Exp. Botan., **30**, 215 – 223.
- Neubert, A., Kley, D., Wildt, J., Segschneidesdewr, H. J. and Forstel, H. (1993) Uptake of NO,NO<sub>2</sub> & O<sub>3</sub> by sunflower (*Helianthous annoul*) and tobacco plant (*Nicotian tobacum*) and its dependence on stomatal conductivity. Atmos. Environ., **27** (14), 2137 -2145.
- Pratiba, S. and Madhu, S. (2000) Change in chlorophyll and total free amino acids of Gram in response to SO<sub>2</sub> exposure. Lander field conditions Poll. Res., **19** (1), 95-97.
- Panday, D., Panday, V. S. and srivaotava, R. P. (1974) A note on the effect of the size of corms on the sprouting and flowering of saffron. Progr. Horticult., **6** (2-3), 89-99.
- Sivasamy, N. and Srinivasan, V. (1997) Environmental pollution and its control by trees. *Employement news*. New Delhi (29<sup>th</sup> march -4<sup>th</sup> April 1997) xxi. **52**, 1-2.

Tingley, D. T., Reinert, R.A., Dunning, J. A. and Heck, W.W.(1971) Vegetation injury from the interaction of NO<sub>2</sub> & SO<sub>2</sub>. *Phytopathology*, **61**, 5106-5111.

Vorobeichik, E. L. (2002) Changes in spatial structure of the destruction process under the conditions of atmospheric pollution of forest ecosystems. *Izv Akad Nauk Ser Bio.*, **3**, 368-79.

Well burn, A. R., Robinson, D. and Walmsley, C. (1981) Bio Chemical explanation of more than additive

inhibitory effects of low atmosphere levels of SO<sub>2</sub> and NO<sub>2</sub> upon Plants. *New Physiologist*, **88**, 223–237.

West, P.W. and Gaek, G. C. (1956) Fixation of SO<sub>2</sub> as sulfitomercurate(11) and subsequent colorimetric estimation. *Analyt. Chem.*, **28**, 1816-1819.

Yi, H., Liu, J. and Zheng, K. (2005) Effect of sulphur dioxide hydrates on cell cycle, sister chromatid exchange and micronuclei in barley. *Ecotoxic. Environ. Saf.*, **62** (3), 421-6.